

TRONIX[®] 700 SC

An agricultural herbicide for use in sugar beet, fodder beet, red beet and mangels


A suspension concentrate containing 700 g/L metamitron (58.3 % w/w)
MAPP No.: 18137 **For Professional Use Only**

IMPORTANT INFORMATION: FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum individual dose (L product/ha)	Max total dose (L product/ha)	Latest time of application
Sugar beet, fodder beet, red beet, mangels	2	5	Before crop leaves meet between the rows.

Other Specific Restrictions: The minimum interval between applications is 6 days. Fodder beet and mangels must not be grazed by livestock or harvested for animal consumption until at least 32 days following the last application.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



WARNING

H302 Harmful if swallowed.
H410 Very toxic to aquatic life with long lasting effects.

P102 Keep out of reach of children.
P264 Wash hands thoroughly after handling.
P270 Do not eat, drink or smoke when using this product.

P391 Collect spillage.
P501 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

EUH208 Contains 5-chloro-2-methylisothiazol-3-one/2-methylisothiazol-3-one. May produce an allergic reaction.

EUH401 To avoid risks to human health and the environment, comply with the instructions for use.

UK

Content: **5L**e

Important: The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

PROTECT FROM FROST

SHAKE WELL BEFORE USE

Avoid spray drift

24-hour emergency number:
0032 14 58 45 45



For warehouse purposes only

© Registered trademark of Globachem N.V.



Marketing company: **Belchim Crop Protection Ltd**
1b Fenice Court, Phoenix Park, Eaton Socon, St Neots, Cambridgeshire PE19 8EW | www.belchim.co.uk | info-uk@belchim.com

REGISTRATION HOLDER: **GLOBACHEM NV**
Lichtenberglaan 2019 | BE-3800 Sint-Truiden
www.globachem.com



This booklet is part of the approved Product Label.

SAFETY PRECAUTIONS

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.
- However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.
- TAKE OFF IMMEDIATELY all contaminated clothing.
- WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.
- WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE.

ENVIRONMENTAL PROTECTION

- Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.
- Extreme care must be taken to avoid spray drift on to non-crop plants outside of the target area.

STORAGE AND DISPOSAL

- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
- WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.
- DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

TRONIX 700 SC is a versatile crop safe selective herbicide with contact and residual properties acting by both leaf and root uptake for the control of annual weeds in sugar beet, red beet, fodder beet and mangels. TRONIX 700 SC is absorbed by the roots of any emerging weeds.

RESTRICTIONS/WARNINGS

Pre-emergence application is not recommended on organic soils.

Factors affecting crop tolerance

TRONIX 700 SC exhibits great safety to beet combined with consistent weed control when used as recommended. The beet plant is, however, very sensitive to herbicide treatments if the following stress factors occur: previously applied pesticides and herbicides (other than TRONIX 700 SC pre-emergence on mineral soils), soft growth after prolonged rain, sudden changes in temperature, high light intensity, high radiant temperatures, prolonged low temperature periods, insect and fungal attack, damage by wind blows, nutrient deficiencies such as manganese. If these or any other stress-inducing conditions occur, some cotyledon tip scorch may occasionally result. Under these conditions crop tolerance to TRONIX 700 SC + CROPSPRAY 11E may be reduced, particularly when used in conjunction with other herbicides either as tank-mixes or sequentially.

Uneven application

Spraying can be uneven due to the movement of spraying equipment, ground undulations, proximity of hedgerows etc. This can lead to local high dosages of chemical and damage to the crop for which no responsibility can be accepted by the suppliers.

WEEDS CONTROLLED

Pre-emergence followed by post-emergence applications

Weeds controlled from a pre-emergence application of TRONIX 700 SC followed by a program of well timed sprays of post-emergence (at the early cotyledon stage of the weeds) applications of TRONIX 700 SC + CROPSPRAY 11E.

Susceptible

Dead-nettle, red	Groundsel
Fat-hen	Scarlet pimpernel

Moderately susceptible

Chickweed, common

Resistant

Black-bindweed	Knotgrass
Cleavers	Pansy, field

Post-emergence programme

Weeds controlled at early cotyledon stage from a series of well timed sprays of TRONIX 700 SC + CROPSPRAY 11E applications

Susceptible

Annual meadowgrass	Chickweed, common
Dead-nettle, red	Fat-hen
Groundsel	Hemp-nettle, common

Nettle, small	Orache, common
Pineappleweed	Poppy, common
Shepherd's purse	Speedwells

Moderately susceptible

Forget-me-not, field	Knotgrass
Mayweeds	Pansy, field
Penny-cress, field	Persicaria, pale
Redshank	Scarlet pimpernel

Moderately resistant

Fumitory, common	Parsley, fool's
------------------	-----------------

Resistant

Black-bindweed	Cleavers
Perennial weeds	

It is important to adhere to the full programme of sprays to ensure complete kill, particularly under dry soil conditions.

TRONIX 700 SC is recommended for use in all varieties of sugar beet grown on mineral and organic soils and red beet, fodder beet and mangels grown on mineral soils.

CROP SPECIFIC INFORMATION

TRONIX 700 SC is recommended for use in all varieties of sugar beet grown on mineral and organic soils and red beet, fodder beet and mangels grown on mineral soils.

TIME OF APPLICATION

TRONIX 700 SC may be applied either as a pre-emergence application followed by post-emergence applications or as a post-emergence programme.

Pre-emergence of beet followed by post-emergence applications:

Apply the first application of TRONIX 700 SC at the pre-emergence stage of the crop. Prolonged dry weather after application may reduce effectiveness.

Best results will be obtained on a fine, well consolidated seedbed, free from clods and established weeds. Cloddy or fluffy seedbeds or very dry conditions will reduce activity. A second application should be applied post-emergence of the crop at the first weed flush,

when the weeds are at the expanded cotyledon stage. A further application should be applied when the next flush of weeds germinate. This is usually 10-21 days after the first post-emergence application.

Post-emergence programme:

The first application should be made at the cotyledon stage of the earliest germinating weeds. The size of the beet does not matter provided the crop is not under stress. The treatment should be repeated as each flush of weeds reaches the cotyledon stage until weed germination ceases. If weeds have survived the previous spray 7-10 days after treatment, another application should be made even if no new weeds have germinated during the period.

RATE OF USE

Pre-emergence of beet followed by post-emergence applications:

On all mineral soils apply TRONIX 700 SC at 1.65 L/ha in 80-200 litres of water per hectare as an overall pre-emergence application. This should precede two post-emergence sprays of TRONIX 700 SC at 1.65 L/ha + CROPSPRAY 11E at 1.65 L/ha applied in 80-100 L/ha water. Alternatively an application of TRONIX 700 SC may be used pre-emergence at 2.0 L/ha followed by 2 post-emergence applications of TRONIX 700 SC at 1.5 L/ha + CROPSPRAY 11E at 1.65 L/ha.

Post-emergence programme:

An overall programme of three sprays using: TRONIX 700 SC 1.65 L/ha + CROPSPRAY 11E at 1.65 L/ha in 80-100 litres of water per hectare. N.B. The residual activity of TRONIX 700 SC is dependent on the cumulative rate of TRONIX 700 SC applied; therefore, if the full programme of sprays is not used, residual activity and efficacy will be reduced.

FOLLOWING CROPS

Beet crops may be sown at any time following the use of TRONIX 700 SC. Providing 16 weeks elapse from the last application of TRONIX 700 SC, winter cereals may be sown in the same season. Any spring crop may be sown in the season following use of TRONIX 700 SC or TRONIX 700 SC/CROPSPRAY 11E. Mouldboard ploughing to a depth of 15 cm followed by thorough cultivation is recommended before planting any crop.

MIXING AND SPRAYING

Thoroughly shake the pack before use. Add the required quantity of TRONIX 700 SC to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing. When tank-mixes are to be used, each product should be added separately to the tank; the TRONIX 700 SC should be dispersed first unless otherwise specified under the Compatibility section.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely. Pre-emergence of the crop use a FINE-MEDIUM spray and post-emergence a FINE spray as defined by the BCPC system. Avoid spray drift. Do not use finer than 80 mesh filters in spray lines or nozzles.

Clean spray equipment thoroughly after use.

COMPATIBILITY

For information on tank mixes please consult your supplier.

WEED RESISTANCE

This product contains metamilon, a Group C1 herbicide, based on the mode of action classification system of the Herbicide Resistance Action Committee. Repeated use of herbicides with the same mode of action can increase the risk of strains of weed developing resistance to these compounds, leading to poor control. In order to minimize the risk, a strategy for preventing and managing such resistance should be adopted. Growers are advised to apply products containing herbicides with different modes of action in sequence or tank mix where two or more components are active against the target weeds. Use the recommended rate of TRONIX 700 SC and the correct application timing for the hardest to control weed species present in the field. The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling. Further guidance on weed resistance management is available from the Herbicide Resistance Action Group (HRAC) and Weed Resistance Action Group (WRAG). Follow WRAG guidelines.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

Other brand names referred to on this label are trademarks of other manufacturers in which proprietary rights may exist.

To access the Safety Data Sheet for this product, scan the QR code or use the weblink below:



www.belchim.co.uk/pdf/MSDS/Tronix_MSDS.pdf

Alternatively, contact your supplier.

For warehouse purposes only